

The Southwest Miramichi Smolt Study 2007

Miramichi Salmon Association
Department of Fisheries and Oceans

By

Jenny Reid
Biologist

November 21, 2007

Introduction

Over the past three decades, there has been a continuing and recognizable need for conservation efforts to sustain Atlantic salmon stocks in the Miramichi River. Over that time, despite major management actions such as the closing of commercial fisheries in both the Maritimes and Newfoundland, the annual returns have fallen below expectations. In very recent years, minimum spawning requirements for Atlantic salmon have not been realized in the Miramichi River system. There clearly is a need for further management action to find the cause of declining stocks and to provide a solution, which will bring a recovery to them. One of the large questions is to determine whether the problem is merely in the ocean, or in the river, or perhaps both.

Although electrofishing studies have consistently indicated an abundance of juvenile salmon (fry and parr) in the river, the adult returns have been much lower than should logically be expected. It has been assumed that smolt (young salmon ready to migrate to the ocean) production would be consistent with the high levels of juveniles, but recent DFO surveys on the Northwest Miramichi indicate that this assumption may be incorrect. Hence, an accurate estimation of the total smolt population migrating from the Miramichi River is an essential component to understanding and managing the regeneration process of the Atlantic salmon in this watershed.

Methods

The method used to obtain the smolt inventory estimates was a mark and recapture concept. On the Cains and Dungarvon Rivers, rotary screw traps (RST) or smolt wheels were used to capture smolts for tagging. The Cains smolt wheel operated from May 4-June 5, and the Dungarvon smolt wheel operated from May 3-June 6, 2007. The smolt wheel was strung across the river by an overhead cable and floats on the top of the water by two large pontoons. The current forced the partially submerged wheel to rotate. Any fish that entered the trap were guided into the trap's holding box which is located at the back of the smolt wheel. The rotating wheel prevented the fish from swimming out of the trap. All the fish in the live-box were collected and sorted. Each species caught was identified, counted and released, except for salmon smolts, which were measured for fork length and then tagged with streamer research tags. Scale samples were also taken from up to five smolts per day for age analysis. After the smolts were tagged they were moved upstream of the smolt wheel. The percentage of tagged smolts that are recaptured at the smolt wheel allow us to estimate the number of smolts moving out of that particular tributary.

A single large trapnet was installed in the estuary of the Southwest Miramichi at Millerton to capture smolts moving from freshwater into the estuary from May 12 to June 11, 2006. Tagged smolts captured at the Millerton trap net allow us to get an estimate of the smolts moving out of the Southwest Miramichi. The Millerton trapnet efficiency is calculated by the percentage of these tagged smolts that are recaptured, and this trap efficiency is then extrapolated to estimate the total smolt run from the number of untagged smolts also captured there. This latter facility was fished daily, generally at low

tide, and the smolts were sorted from the rest of the species captured. Each day, subsamples of up to 100 smolts were measured and 20 were sampled in detail for length, weight, sex and age. All smolts captured were counted and checked for missing adipose fin clips and tags.

Results

The peak of the smolt run occurred on May 21st for both the Cains and Dungarvon Rivers. This year we tagged 859 smolts on the Cains and 2718 smolts on the Dungarvon River. We estimated that 70,000 smolts moved from the Dungarvon River and 56,000 from the Cains River downstream to the ocean in the spring of 2007. The smolt production on the Dungarvon River in 2007 was the highest ever recorded in the past 6 years, 48,000, exceeding the target production of 3.0 smolts per 100m² of habitat (Table 1). Of the tagged fish 27 smolts from the Cains and 58 from the Dungarvon River were later recaptured at the trap net near the estuary at Millerton. We estimated a total smolt production of 1.34 million smolts in the Southwest Miramichi, the highest in the past five years. In addition, 0.5% of the Cains, 7.7% of the Dungarvon and 0.5% of the Southwest Miramichi smolt runs were comprised of salmon smolts with clipped adipose fins which were stocked by MSA a few years earlier.

Table 1. Estimate for salmon smolt movements (using 0% mortality factor) for SW Miramichi 2001 – 2007, Cains River 2002 – 2007 and Dungarvon River for 2002 - 2007. The desired target for smolt production on the Miramichi River is 3.0 smolts/100m² of habitat. *Note: This is only a minimum smolt estimate as wheel was not operational for three days during typical smolt peak time.

River	Year	Smolt estimate	Smolts/100m²
SW Miramichi	2001	500,000	1.4
	2002	637,000	1.8
	2003	534,400	1.5
	2004	1,152,000	3.3
	2005	N/A	N/A
	2006	1,336,000	3.8
	2007	1,340,000	3.8
Cains	2002	38,500	0.8
	2003	47,000	1.0
	2004	78,000	1.7
	2005	N/A	N/A
	2006	98,000	2.1
	2007	56,000	1.2
Dungarvon	2002	43,300	1.8
	2003	34,700	1.6
	2004	38,000	1.7
	2005	58,000*	2.6
	2006	48,000	2.1
	2007	70,000	3.2